

**Table 7.24A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2007 and 2008**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	10,870	11,087
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3,215	3,406	4,836	5,264
Cigars	7,316	7,505	5,947	5,621
Pipe Tobacco	1,138	957	907	920
<b>ALCOHOL</b>	40,166	40,290	86,594	88,684
Binge Alcohol Use <sup>2</sup>	27,031	26,675	30,747	31,421
Heavy Alcohol Use <sup>2</sup>	9,877	10,032	7,134	7,259

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.24B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2007 and 2008**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	5.8	5.8
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	5.4	5.7	2.6	2.8
Cigars	12.2	12.6	3.2	3.0
Pipe Tobacco	1.9	1.6	0.5	0.5
<b>ALCOHOL</b>	66.9	67.4	46.1	46.7
Binge Alcohol Use <sup>2</sup>	45.0	44.6	16.4	16.5
Heavy Alcohol Use <sup>2</sup>	16.4	16.8	3.8	3.8

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.25A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2007 and 2008**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	646	583
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	324	292	293	255
Cigars	699 <sup>a</sup>	598	371	345
Pipe Tobacco	135	128	46	49
<b>ALCOHOL</b>	1,512 <sup>a</sup>	1,329	2,509 <sup>a</sup>	2,306
Binge Alcohol Use <sup>2</sup>	1,220 <sup>a</sup>	1,058	1,227	1,130
Heavy Alcohol Use <sup>2</sup>	422 <sup>b</sup>	319	169	172

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.25B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2007 and 2008**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	2.8	2.6
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	13.0	12.9	1.3	1.1
Cigars	28.2	26.5	1.6	1.5
Pipe Tobacco	5.5	5.6	0.2	0.2
<b>ALCOHOL</b>	60.9	58.8	11.0 <sup>a</sup>	10.2
Binge Alcohol Use <sup>2</sup>	49.2	46.8	5.4	5.0
Heavy Alcohol Use <sup>2</sup>	17.0	14.1	0.7	0.8

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.26A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2007 and 2008**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	1,859	1,871
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,125	1,105	596	667
Cigars	2,554	2,485	1,319	1,236
Pipe Tobacco	279	302	115	155
<b>ALCOHOL</b>	9,361	9,325	10,673	10,824
Binge Alcohol Use <sup>2</sup>	7,446	7,488	6,228	6,022
Heavy Alcohol Use <sup>2</sup>	3,140	3,080	1,680	1,681

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.26B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2007 and 2008**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	8.9	8.8
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	9.5	9.4	2.9	3.2
Cigars	21.6	21.1	6.3	5.8
Pipe Tobacco	2.4	2.6	0.6	0.7
<b>ALCOHOL</b>	79.1	79.3	51.1	51.1
Binge Alcohol Use <sup>2</sup>	62.9	63.7	29.8	28.4
Heavy Alcohol Use <sup>2</sup>	26.5	26.2	8.0	7.9

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.27A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2007 and 2008**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	8,365	8,633
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,766	2,010	3,947	4,341
Cigars	4,064	4,422	4,256	4,040
Pipe Tobacco	724	528	746	717
<b>ALCOHOL</b>	29,293	29,636	73,412	75,555
Binge Alcohol Use <sup>2</sup>	18,364	18,129	23,292	24,268
Heavy Alcohol Use <sup>2</sup>	6,315	6,633	5,285	5,407

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.27B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2007 and 2008**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	5.8	5.9
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3.9	4.4	2.7	3.0
Cigars	8.9	9.7	3.0	2.8
Pipe Tobacco	1.6	1.2	0.5	0.5
<b>ALCOHOL</b>	64.0	64.8	50.9	51.7
Binge Alcohol Use <sup>2</sup>	40.1	39.6	16.2	16.6
Heavy Alcohol Use <sup>2</sup>	13.8	14.5	3.7	3.7

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.